

result in loss of user privileges and other penalties.

\*\*\*\*\* STN Columbus \*\*\*\*\*

FILE 'HOME' ENTERED AT 14:15:02 ON 09 FEB 2007

=> file reg

COST IN U.S. DOLLARS

SINCE FILE

ENTRY

TOTAL

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 14:15:16 ON 09 FEB 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 8 FEB 2007 HIGHEST RN 920112-67-0

DICTIONARY FILE UPDATES: 8 FEB 2007 HIGHEST RN 920112-67-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 30, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=> ....Testing the current file.... screen


ENTER SCREEN EXPRESSION OR (END):end

=> screen 1006

L1 SCREEN CREATED

=>

Uploading C:\Program Files\Stnexp\Queries\10560764.str



```

chain nodes :
7 8 9 10 11 12 13 20 21 22 23 24 25 26 27
ring nodes :
1 2 3 4 5 6 14 15 16 17 18 19
chain bonds :
6-7 7-8 8-9 9-10 10-11 10-13 11-12 12-14 17-20 20-21 21-22 21-25 22-23
22-24 25-26 26-27
ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6 14-15 14-19 15-16 16-17 17-18 18-19
exact/norm bonds :
9-10 10-13 12-14 21-25 22-23 22-24
exact bonds :
6-7 7-8 8-9 10-11 11-12 17-20 20-21 21-22 25-26 26-27
normalized bonds :
1-2 1-6 2-3 3-4 4-5 5-6 14-15 14-19 15-16 16-17 17-18 18-19

```

```

Match level :
1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:Atom 8:Atom 9:Atom
10:Atom 11:Atom 12:Atom 13:Atom 14:CLASS 15:CLASS 16:CLASS 17:CLASS
18:CLASS 19:CLASS 20:Atom 21:Atom 22:Atom 23:Atom 24:Atom 25:Atom 26:Atom
27:Atom

```

L2 STRUCTURE UPLOADED

=> que L2 AND L1

L3 QUE L2 AND L1

=> d L1


L1 HAS NO ANSWERS

L1 SCR 1006

=> d L2

L2 HAS NO ANSWERS

L2 STR



Structure attributes must be viewed using STN Express query preparation.

=> s L2 full  
FULL SEARCH INITIATED 14:15:58 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 716 TO ITERATE

100.0% PROCESSED 716 ITERATIONS 19 ANSWERS  
SEARCH TIME: 00.00.01

L4 19 SEA SSS FUL L2

=> file caplus  
COST IN U.S. DOLLARS SINCE FILE TOTAL  
ENTRY SESSION  
FULL ESTIMATED COST 172.10 172.31

FILE 'CAPLUS' ENTERED AT 14:16:03 ON 09 FEB 2007  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 9 Feb 2007 VOL 146 ISS 8  
FILE LAST UPDATED: 8 Feb 2007 (20070208/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s L4  
L5 9 L4



=> d L5 1-9 bib abs

L5 ANSWER 1 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2007:61504 CAPLUS  
DN 146:142376  
TI Preparation of phenylpropionic acid derivatives and pharmaceutical compositions thereof  
IN Bjoerk, Seth  
PA Astrazeneca AB, Swed.  
SO PCT Int. Appl., 57pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2007008156	A1	20070118	WO 2006-SE864	20060710
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG,			

US, UZ, VC, VN, ZA, ZM, ZW  
 RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,  
 IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ,  
 CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH,  
 GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,  
 KG, KZ, MD, RU, TJ, TM

PRAI SE 2005-1644 A 20050711  
 GI




AB The title phenylpropionic acid derivs. I [wherein n = 1-2; R1 = H, Cl, CF3, or OCF3; R2 = H or F; R3 = alkyl] or tert-butylamine salts thereof were prepared as PPAR active compds. for treatment of metabolic syndrome including type 2 diabetes mellitus (no data). For example, II and II•tert-butylamine were prepared in a multi-step synthesis. Pharmaceutical compns. were described.

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 2 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 2006:605020 CAPLUS  
 DN 145:83115  
 TI Preparation of tris(hydroxymethyl)methylamine and ethanolamine salts of (2S)-2-ethoxy-3-(4-{2-[hexyl(2-phenylethyl)amino]-2-oxoethoxy}phenyl)propanoic acid for treating lipid disorders  
 IN Booth, Rebecca J.; Dahlstroem, Mikael  
 PA AstraZeneca AB, Swed.  
 SO PCT Int. Appl., 39 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006065214	A1	20060622	WO 2005-SE1916	20051214
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,				

KG, KZ, MD, RU, TJ, TM  
 PRAI SE 2004-3072 A 20041216  
 GI



I


AB The invention relates to a compound selected from one or more of the following: a tris(hydroxymethyl)methylamine salt or an ethanolamine salt of title compound I or a pharmaceutical composition comprising the compound  
 Thus I was prepared in 4 steps from Et (S)-2-ethoxy-3-(4-hydroxyphenyl)propanoate, benzyl bromoacetate, and N-hexyl-2-phenylethylamine. X-ray powder diffraction patterns for bot salts of I are given: Both salts have an EC50 of less than 0.5  $\mu\text{mol/l}$  for PPAR $\alpha$ .

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 3 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 2005:1335635 CAPLUS  
 DN 144:69628  
 TI Preparation of phenoxyacetamide derivatives as modulators of peroxisome proliferator-activated receptors (PPAR)  
 IN Alstermark, Eva-Lotte Lindstedt; Olsson, Anna Christina; Li, Lanna  
 PA Swed.  
 SO U.S. Pat. Appl. Publ., 47 pp., Cont.-in-part of U.S. Ser. No. 499,261.  
 CODEN: USXXCO  
 DT Patent  
 LA English  
 FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2005282822	A1	20051222	US 2004-26806	20041230
	WO 2003051821	A1	20030626	WO 2002-GB5738	20021218
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	WO 2003051822	A1	20030626	WO 2002-GB5744	20021218
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CN 1896045	A	20070117	CN 2006-10007173	20021218
	WO 2004056748	A1	20040708	WO 2003-GB5602	20031219
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,				

CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,  
 GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
 LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO,  
 NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ,  
 TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,  
 BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
 ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,  
 TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG  
 WO 2004113270 A2 20041229 WO 2004-EP6597 20040617  
 WO 2004113270 A3 20050331  
 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,  
 CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,  
 GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
 LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI,  
 NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,  
 TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,  
 EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE,  
 SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,  
 SN, TD, TG  
 EP 1676833 A1 20060705 EP 2006-5766 20040617  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR  
 JP 2005336209 A 20051208 JP 2005-235794 20050816  
 JP 2006045240 A 20060216 JP 2005-253346 20050901  
 JP 2006298924 A 20061102 JP 2006-123399 20060427  
 JP 2006298925 A 20061102 JP 2006-139673 20060519  
 PRAI SE 2001-4334 A 20011219  
 WO 2002-GB5738 W 20021218  
 WO 2002-GB5744 A 20021218  
 GB 2002-29931 A 20021221  
 GB 2003-14079 A 20030618  
 WO 2003-GB305602 A 20031219  
 WO 2004-EP6597 A 20040617  
 US 2005-499261 A2 20050304  
 CN 2002-828123 A3 20021218  
 JP 2003-552709 A3 20021218  
 JP 2003-552710 A3 20021218  
 JP 2004-561668 A3 20031219  
 EP 2004-740044 A3 20040617  
 JP 2006-515989 A3 20040617  
 OS MARPAT 144:69628  
 GI



AB The phenyl-, phenoxy-, or phenylthioalkanamidetitle compds., (in particular phenoxyacetamide derivs.) (I) [A is situated in the ortho, meta or para position and represents CR<sub>3</sub>R<sub>4</sub>CR<sub>1</sub>R<sub>2</sub>COR, CR<sub>3</sub>:CR<sub>1</sub>COR (wherein R = H, alkyl, (un)substituted HO or NH<sub>2</sub>; R<sub>1</sub> = alkyl, aryl, alkenyl, alkynyl, or when A is CR<sub>3</sub>R<sub>4</sub>CR<sub>1</sub>R<sub>2</sub>COR, R<sub>1</sub> can also be cyano, (un)substituted HO, SH, OCONH<sub>2</sub>, SO<sub>2</sub>NH<sub>2</sub>, CO<sub>2</sub>H, etc.; R<sub>2</sub> = H, halogen, alkyl, aryl, alkylaryl; R<sub>3</sub>, R<sub>4</sub> = H, alkyl, aryl, alkylaryl); Y = O, S, a single bond; n = an integer of 1-4; X = alkyl; R<sub>5</sub>, R<sub>6</sub> = H, each (un)substituted C1-13 alkyl, C2-10 alkenyl, or C2-10 alkynyl; or R<sub>5</sub>, R<sub>6</sub> = each (un)substituted C3-8

cycloalkyl, C3-C8 cycloalkenyl, aryl, heterocyclyl, or heteroaryl; or R5 and R6 together with the nitrogen atom to which they are attached form a single or a fused heterocyclic system] are prepared These compds. are useful in treating clin. conditions including lipid disorders (dyslipidemias) whether or not associated with insulin resistance, and other manifestations of the metabolic syndrome. Thus, a solution of 0.598 g N-butyl-N-[2-fluoro-4-(trifluoromethyl)benzyl]amine and 0.593 g [4-((2S)-2,3-diethoxy-3-oxopropyl)phenoxy]acetic acid in 20 mL CH<sub>2</sub>Cl<sub>2</sub> was treated with 0.80 mL N,N-diisopropylethylamine and 0.674 g O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate and the reaction mixture was stirred at room temperature overnight to give, after workup and silica gel chromatog., 74% Et (2S)-3-[4-[2-[butyl[2-fluoro-4-(trifluoromethyl)benzyl]amino]-2-oxoethoxy]phenyl]-2-ethoxypropanoate (II). A solution of 0.748 g II in 70 mL MeCN was treated with 35 mL 0.10 M LiOH and the reaction mixture was stirred at room temperature overnight, neutralized with 5% HCl, concentrated, acidified with 5% HCl, and extracted



with

EtOAc to give 97% (2S)-3-[4-[2-[butyl[2-fluoro-4-(trifluoromethyl)benzyl]amino]-2-oxoethoxy]phenyl]-2-ethoxypropanoic acid (III). III showed EC<sub>50</sub> of 0.001 µmol/L for human PPAR $\alpha$ .

L5 ANSWER 4 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 2004:1154649 CAPLUS  
 DN 142:93514  
 TI Preparation of phenylpropanoic acid derivatives as PPAR $\alpha$  agonists  
 IN Li, Lanna; Lindstedt-Alstermark, Eva-Lotte; Olsson, Christina  
 PA Astrazeneca Ab, Swed.  
 SO PCT Int. Appl., 100 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004113270	A2	20041229	WO 2004-EP6597	20040617
	WO 2004113270	A3	20050331		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2004249409	A1	20041229	AU 2004-249409	20040617
	CA 2528234	A1	20041229	CA 2004-2528234	20040617
	US 2005148656	A1	20050707	US 2003-518777	20040617
	EP 1675820	A2	20060705	EP 2004-740044	20040617
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
	EP 1676833	A1	20060705	EP 2006-5766	20040617
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
	BR 2004011484	A	20060725	BR 2004-11484	20040617
	CN 1835913	A	20060920	CN 2004-80023304	20040617
	JP 2006527730	T	20061207	JP 2006-515989	20040617
	US 2005282822	A1	20051222	US 2004-26806	20041230
	NO 2005005892	A	20060106	NO 2005-5892	20051212
	JP 2006298925	A	20061102	JP 2006-139673	20060519
	US 2006258866	A1	20061116	US 2006-477168	20060628
PRAI	GB 2003-14079	A	20030618		
	SE 2001-4334	A	20011219		

OS MARPAT 142:93514  
GI



AB Title compds. represented by the formula I [wherein A = CR3(R4)CR1(R2)COR or C(R3):C(R1)COR; R = H, alkoxy, (alkyl)aryloxy, amino, etc.; R1 = alkyl, aryl, alkenyl, alkynyl, etc.; R2 = H, halo, alkyl, (alkyl)aryl; R3, R4 = independently H, alkyl, (alkyl)aryl; T = O, S or a single bond; n = 1-4; R5, R6 = independently selected substituent comprising C, H, N, O, S, Se, P or halo; with provisos; optical isomers and racemates thereof as well as pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof] were prepared as PPAR $\alpha$  agonists. For example, II was given in a multi-step synthesis starting from the reaction of 2,4-difluorobenzylamine with octanoic acid. I had EC50 values of less than 0.1  $\mu$ mil/L for PPAR $\alpha$  and showed the ration of the EC50(PPAR $\gamma$ ) with EC50(PPAR $\alpha$ ) is greater than 150:1. Thus, I and their pharmaceutical compns. are useful for the treatment of clin. conditions including lipid disorders (dyslipidemias) whether or not associated with insulin resistance (no data).

L5 ANSWER 5 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2004:1127321 CAPLUS  
DN 142:49239

TI    Pharmaceutically useful salts (2S)-2-ethoxy-3-(4-{2[hexyl(2-phenylethyl)amino]-2-oxoethoxy}phenyl)propanoic acid, preparation thereof, and therapeutic use

IN Ragnar, Ralf; Stahle, Erica

PA Astrazeneca AB, Swed.

SO PCT Int. Appl., 38 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004110985	A1	20041223	WO 2004-SE965	20040616



W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

AU 2004247611 A1 20041223 AU 2004-247611 20040616  
 CA 2527608 A1 20041223 CA 2004-2527608 20040616  
 EP 1638921 A1 20060329 EP 2004-736956 20040616

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR

BR 2004011455 A 20060718 BR 2004-11455 20040616  
 CN 1805922 A 20060719 CN 2004-80016838 20040616  
 JP 3836498 B2 20061025 JP 2006-517040 20040616  
 JP 2006527767 T 20061207  
 US 2006194879 A1 20060831 US 2005-560127 20051209  
 NO 2005005923 A 20060106 NO 2005-5923 20051213

PRAI GB 2003-14136 A 20030618  
 WO 2004-SE965 W 20040616

AB The invention discloses a calcium or magnesium salt of (2S)-2-ethoxy-3-(4-{2[hexyl(2-phenylethyl)amino]-2-oxoethoxy}phenyl)propanoic acid. Comps. of the invention (preparation included) may be used to treat e.g. dyslipidemia and type 2 diabetes.

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 6 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 2004:1127320 CAPLUS  
 DN 142:49238  
 TI Pharmaceutically useful salts of (2S)-2-ethoxy-3-[4-(2-(hexyl(2-phenylethyl)amino)-2-oxoethoxy)phenyl]propanoic acid, their preparation, and their therapeutic use  
 IN Aurell, Carl-Johan; Dahlstroem, Mikael; Lindstedt-Alstermark, Eva-Lotte; Minidis, Anna; Ohlsson, Bengt; Stahle, Erica  
 PA Astrazeneca AB, Swed.  
 SO PCT Int. Appl., 47 pp.  
 CODEN: PIXXD2

DT Patent  
 LA English

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004110984	A1	20041223	WO 2004-SE964	20040616
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2004247610	A1	20041223	AU 2004-247610	20040616
CA 2528932	A1	20041223	CA 2004-2528932	20040616
EP 1638922	A1	20060329	EP 2004-749009	20040616
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
CN 1809529	A	20060726	CN 2004-80016948	20040616


BR 2004011525	A	20060801	BR 2004-11525	20040616
JP 3822900	B2	20060920	JP 2006-517039	20040616
JP 2006527766	T	20061207		
NO 2005005922	A	20060106	NO 2005-5922	20051213
US 2006142389	A1	20060629	US 2005-560657	20051213
PRAI GB 2003-14129	A	20030618		
WO 2004-SE964	W	20040616		

AB The invention discloses salts of (2S)-2-ethoxy-3-[4-(2-(hexyl(2-phenylethyl)amino)-2-oxoethoxy)phenyl]propanoic acid e.g. the L-arginine salt. Preparation of compds. of the invention is described. The compds. of the invention are useful in the treatment of e.g. dyslipidemias and other manifestations of the metabolic syndrome.

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 7 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2004:1127318 CAPLUS  
DN 142:56001  
TI Preparation of (2S)-3-(4-{2-[amino]-2-oxoethoxy}phenyl)-2-ethoxypropanoic acid derivatives  
IN Aurell, Carl-Johan; Macedo, Emmanuel; Minidis, Anna; Yousefi-Salakdeh, Esmail  
PA Astrazeneca Ab, Swed.  
SO PCT Int. Appl., 16 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004110982	A1	20041223	WO 2004-SE966	20040616
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2004247612	A1	20041223	AU 2004-247612	20040616
	CA 2528933	A1	20041223	CA 2004-2528933	20040616
	EP 1638920	A1	20060329	EP 2004-736958	20040616
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, HR				
	CN 1809528	A	20060726	CN 2004-80017131	20040616
	BR 2004011558	A	20060801	BR 2004-11558	20040616
	JP 3822901	B2	20060920	JP 2006-517041	20040616
	JP 2006527768	T	20061207		
	NO 2005005924	A	20060105	NO 2005-5924	20051213
	US 2006142392	A1	20060629	US 2005-560764	20051213
PRAI	GB 2003-14134	A	20030618		
	WO 2004-SE966	W	20040616		
OS	MARPAT 142:56001				
GI					



I

AB The present invention provides a process for preparation of the title compds. I (R = H, R1 = n-C6H13) by reacting I (R = H, or protecting group, R1 = H) with C6H13X (X = leaving group) in the presence of a base and inert solvent at a temperature in the range -25°C to 150°C and optionally, when OR represents a protecting group, removal of the protecting group.

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 8 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2004:2837 CAPLUS

DN 140:59411

TI Preparation of phenoxyalkanamides as amide linker peroxisome proliferator activated receptor agonists for treating and/or preventing diabetes mellitus and syndrome X

IN Ferritto Crespo, Rafael; Martin, Jose Alfredo; Martin-Ortega, Finger Maria Dolores; Rojo Garcia, Isabel; Shen, Quanrong; Warshawsky, Alan M.; Xu, Yanping

PA Eli Lilly and Company, USA

SO PCT Int. Appl., 168 pp.



CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004000789	A1	20031231	WO 2003-US16207	20030611
	WO 2004000789	A9	20040311		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2488972	A1	20031231	CA 2003-2488972	20030611
	AU 2003241579	A1	20040106	AU 2003-241579	20030611
	EP 1517882	A1	20050330	EP 2003-731326	20030611
	R:				
	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	BR 2003011834	A	20050412	BR 2003-11834	20030611
	CN 1662487	A	20050831	CN 2003-814173	20030611
	JP 2005529975	T	20051006	JP 2004-515700	20030611
	US 2006111406	A1	20060525	US 2004-517581	20041208
PRAI	US 2002-390102P	P	20020619		
	WO 2003-US16207	W	20030611		
OS	MARPAT 140:59411				
GI					



AB The present invention is directed to phenoxyalkyl amides (shown as I; variables defined below; e.g. II), compns., and their use as peroxisome proliferator activated receptor agonists for treating and/or preventing diabetes mellitus and syndrome X. The binding and cotransfection efficacy values found for compds. of this invention that are useful for modulating a PPAR $\alpha$  receptor are about <100 nM and >50%, resp. Although the methods of preparation are not claimed, .apprx.140 example preps. of I are included. For example, II was prepared in 3 steps starting from (2S)-2-ethoxy-3-(4-hydroxyphenyl)propionic acid Me ester, (2S)-2-hydroxypropionic acid benzyl ester and involving intermediates (2S)-3-[4-[[[(1R)-1-[(benzyloxy)carbonyl]ethyl]oxy]phenyl]-2-ethoxypropionic acid Et ester and (2S)-3-[4-[[[(1R)-1-carboxyethyl]oxy]phenyl]-2-ethoxypropionic acid. For I: R1 = H, C1-C8 alkyl, C3-C6 cycloalkyl, aryl-C0-4-alkyl, heteroaryl-C0-4-alkyl, aminoC1-C4alkyl, C3-C6 cycloalkylaryl-C0-2-alkyl, arylheteroC1-C8alkyl, -CHC(O)C1-C4 alkoxy, C0-4-alkyl-C(O)heteroC1-C8alkyl, and -CH2C(O)-R15R16. R2 = C1-C8 alkyl, C3-C6 cycloalkyl, aryl-C0-C4-alkyl, heteroaryl-C0-C4-alkyl, heteroC1-C6cycloalkylaryl, heteroC1-C6cycloalkylarylC1-C4alkyl, aminoC1-C4alkyl, C3-C6 cycloalkylaryl-C0-C2-alkyl, arylheteroC1-C8alkyl, C0-C4-alkyl-C(O)heteroC1-C8alkyl, -CH(C(O)OCH3)benzyl, and -CH2C(O)R15''R16''. R1 and R2 together may form a heterocyclic ring which heterocyclic ring is (un)substituted with 1-3 substituents R1' and which heterocyclic ring is optionally fused with an aryl; E = C(R3)(R4)A, (CH2)nCOOR13, aryl-C0-C4-alkyl, thio-C1-C4-alkyl, thioaryl, arylC1-C4alkoxy, C1-C4alkoxy C1-C4alkyl, aminoaryl, and aminoC1-C4alkyl. R5 and R6 = H, C1-C8 alkyl, aryl-C0-C4-alkyl, heteroaryl-C0-C4-alkyl, C3-C6 cycloalkyl, aryl-C0-C2-alkyl, C3-C6 cycloalkyl-C0-2-alkyl, and -CH2C(O)R17R18.

RE.CNT 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 9 OF 9 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2003:491168 CAPLUS  
DN 139:69049  
TI Preparation of substituted phenylpropionic acid derivatives as agonists to human peroxisome proliferator-activated receptor alpha (PPAR)  
IN Alstermark Lindstedt, Eva-Lotte; Olsson, Anna Christina; Li, Lanna  
PA Astrazeneca AB, Swed.; Astrazeneca UK Limited  
SO PCT Int. Appl., 40 pp.  
CODEN: PIXXD2

DT Patent  
LA English


FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003051821	A1	20030626	WO 2002-GB5738	20021218
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,				

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

CA 2470491	A1	20030626	CA 2002-2470491	20021218
AU 2002366315	A1	20030630	AU 2002-366315	20021218
EP 1458673	A1	20040922	EP 2002-804964	20021218
EP 1458673	B1	20060906		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
BR 2002014988	A	20041214	BR 2002-14988	20021218
HU 200402133	A2	20050228	HU 2004-2133	20021218
CN 1620422	A	20050525	CN 2002-828123	20021218
CN 1620423	A	20050525	CN 2002-828155	20021218
US 2005171204	A1	20050804	US 2003-499261	20021218
JP 2005526011	T	20050902	JP 2003-552709	20021218
JP 3784804	B2	20060614		
TW 253444	B	20060421	TW 2002-91136518	20021218
NZ 533276	A	20060428	NZ 2002-533276	20021218
TW 255807	B	20060601	TW 2002-91136519	20021218
AT 338743	T	20060915	AT 2002-804964	20021218
CN 1896045	A	20070117	CN 2006-10007173	20021218
ZA 2004004657	A	20050829	ZA 2004-4657	20040611
ZA 2004004658	A	20060222	ZA 2004-4658	20040611
NO 2004003023	A	20040715	NO 2004-3023	20040715
US 2005282822	A1	20051222	US 2004-26806	20041230
JP 2005336209	A	20051208	JP 2005-235794	20050816
JP 2006298924	A	20061102	JP 2006-123399	20060427
PRAI SE 2001-4334	A	20011219		
CN 2002-828123	A3	20021218		
JP 2003-552709	A3	20021218		
JP 2003-552710	A3	20021218		
WO 2002-GB5738	W	20021218		
WO 2002-GB5744	A	20021218		
GB 2002-29931	A	20021221		
GB 2003-14079	A	20030618		
WO 2003-GB305602	A	20031219		
WO 2004-EP6597	A	20040617		
US 2005-499261	A2	20050304		
OS MARPAT 139:69049				
GI				



AB The S enantiomer of I, n = 1 or 2, (C<sub>6</sub>H<sub>13</sub> = hexyl) as well as their pharmaceutically acceptable salts, solvates, crystalline forms and prodrugs are synthesized using various solvents and in presence of charcoal-supported palladium catalyst. The utility of these compds. in clin. conditions such as lipid disorders (dyslipidemias) whether or not associated with insulin resistance and therapeutic and other pharmaceutical activities is also investigated. For example, (2S)-3-(4{2-[benzyl(hexyl)amino]-2-oxoethoxy}phenyl)2-ethoxypropionic acid was prepared in 58% yield via reaction of (2S)-2-ethoxy-3-(4-hydroxyphenyl)propanoate and benzyl bromoacetate.

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD

# ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> s (2S)-2-ethoxy-3-(4-{2-oxo-2-[(2-phenylethyl)amino]ethoxy}phenyl)-propanoic acid  
MISSING TERM BEFORE '(2S'

Search expressions cannot begin with operators.

=> s 2-ethoxy-3-(4-{2-oxo-2-[(2-phenylethyl)amino]ethoxy}phenyl)propanoic acid  
MISSING OPERATOR '-ETHOXY-3-(4-{2-OXO-2'

The search profile that was entered contains terms or  
nested terms that are not separated by a logical operator.

=> s phenethylamino ethoxy phenyl propanoic acid derivatives

838 PHENETHYLAMINO  
42317 ETHOXY  
343103 PHENYL  
414 PHENYLS  
343381 PHENYL  
(PHENYL OR PHENYLS)  
1309164 PH  
10070 PHS  
1313504 PH  
(PH OR PHS)  
1566052 PHENYL  
(PHENYL OR PH)  
8991 PROPANOIC  
4311309 ACID  
1568117 ACIDS  
4812460 ACID  
(ACID OR ACIDS)  
340439 DERIVATIVES  
1134482 DERIVS  
1240054 DERIVATIVES  
(DERIVATIVES OR DERIVS)  
L6 0 PHENETHYLAMINO ETHOXY PHENYL PROPANOIC ACID DERIVATIVES  
(PHENETHYLAMINO (W) ETHOXY (W) PHENYL (W) PROPANOIC (W) ACID (W) DERIVATIVES)

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	41.29	213.60
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-7.02	-7.02

STN INTERNATIONAL LOGOFF AT 14:20:59 ON 09 FEB 2007

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssptalxn1621

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 OCT 23 The Derwent World Patents Index suite of databases on STN  
has been enhanced and reloaded  
NEWS 4 OCT 30 CHEMLIST enhanced with new search and display field  
NEWS 5 NOV 03 JAPIO enhanced with IPC 8 features and functionality  
NEWS 6 NOV 10 CA/CAPLUS F-Term thesaurus enhanced  
NEWS 7 NOV 10 STN Express with Discover! free maintenance release Version  
8.01c now available  
NEWS 8 NOV 20 CA/CAPLUS to MARPAT accession number crossover limit increased  
to 50,000  
NEWS 9 DEC 01 CAS REGISTRY updated with new ambiguity codes  
NEWS 10 DEC 11 CAS REGISTRY chemical nomenclature enhanced  
NEWS 11 DEC 14 WPIDS/WPINDEX/WPIX manual codes updated  
NEWS 12 DEC 14 GBFULL and FRFULL enhanced with IPC 8 features and  
functionality  
NEWS 13 DEC 18 CA/CAPLUS pre-1967 chemical substance index entries enhanced  
with preparation role  
NEWS 14 DEC 18 CA/CAPLUS patent kind codes updated  
NEWS 15 DEC 18 MARPAT to CA/CAPLUS accession number crossover limit increased  
to 50,000  
NEWS 16 DEC 18 MEDLINE updated in preparation for 2007 reload  
NEWS 17 DEC 27 CA/CAPLUS enhanced with more pre-1907 records  
NEWS 18 JAN 08 CHEMLIST enhanced with New Zealand Inventory of Chemicals  
NEWS 19 JAN 16 CA/CAPLUS Company Name Thesaurus enhanced and reloaded  
NEWS 20 JAN 16 IPC version 2007.01 thesaurus available on STN  
NEWS 21 JAN 16 WPIDS/WPINDEX/WPIX enhanced with IPC 8 reclassification data  
NEWS 22 JAN 22 CA/CAPLUS updated with revised CAS roles  
NEWS 23 JAN 22 CA/CAPLUS enhanced with patent applications from India  
NEWS 24 JAN 29 PHAR reloaded with new search and display fields  
NEWS 25 JAN 29 CAS Registry Number crossover limit increased to 300,000 in  
multiple databases  
  
NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c, CURRENT  
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.  
  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS LOGIN Welcome Banner and News Items  
NEWS IPC8 For general information regarding STN implementation of IPC 8  
NEWS X25 X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 14:24:08 ON 09 FEB 2007

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'CAPLUS' ENTERED AT 14:24:18 ON 09 FEB 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 9 Feb 2007 VOL 146 ISS 8

FILE LAST UPDATED: 8 Feb 2007 (20070208/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s phenyl propanoic acid derivatives

343103 PHENYL

414 PHENYLS

343381 PHENYL

(PHENYL OR PHENYLS)

1309164 PH

10070 PHS

1313504 PH

(PH OR PHS)

1566052 PHENYL

(PHENYL OR PH)

8991 PROPANOIC

4311309 ACID

1568117 ACIDS

4812460 ACID

(ACID OR ACIDS)

340439 DERIVATIVES

1134482 DERIVS

1240054 DERIVATIVES

(DERIVATIVES OR DERIVS)

L1 7 PHENYL PROPANOIC ACID DERIVATIVES

(PHENYL(W) PROPANOIC(W) ACID(W) DERIVATIVES)

=> d L1 1-7 bib abs

L1 ANSWER 1 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2006:453910 CAPLUS

DN 145:76018



TI Identification of novel PPAR $\alpha$  ligands by the structural modification of a PPAR $\gamma$  ligand

AU Usui, Shinya; Fujieda, Hiroki; Suzuki, Takayoshi; Yoshida, Naoaki; Nakagawa, Hidehiko; Miyata, Naoki

CS Graduate School of Pharmaceutical Sciences, Nagoya City University, Nagoya, Aichi, 467-8603, Japan




SO Bioorganic & Medicinal Chemistry Letters (2006), 16(12), 3249-3254  
 CODEN: BMCLE8; ISSN: 0960-894X  
 PB Elsevier B.V.  
 DT Journal  
 LA English  
 GI



AB To develop novel PPAR $\alpha$  ligands, the authors designed and synthesized several 3-{3-[2-(nonylpyridin-2-ylamino)ethoxy]phenyl} propanoic acid derivs. Compound (I), the meta isomer of a PPAR $\gamma$  agonist (II), has been identified as a PPAR $\alpha$  ligand. The introduction of Me and Et groups at the C-2 position of the propanoic acid of I further improved the PPAR $\alpha$ -binding potency.

RE.CNT 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 2 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 2005:199464 CAPLUS  
 DN 142:430102  
 TI Design, synthesis, and biological activity of novel PPAR $\gamma$  ligands based on rosiglitazone and 15d-PGJ2  
 AU Usui, Shinya; Suzuki, Takayoshi; Hattori, Yoshifumi; Etoh, Kazuma; Fujieda, Hiroki; Nishizuka, Makoto; Imagawa, Masayoshi; Nakagawa, Hidehiko; Kohda, Kohfuku; Miyata, Naoki  
 CS Graduate School of Pharmaceutical Sciences, Nagoya City University, Nagoya, Aichi, 467-8603, Japan  
 SO Bioorganic & Medicinal Chemistry Letters (2005), 15(6), 1547-1551  
 CODEN: BMCLE8; ISSN: 0960-894X  
 PB Elsevier B.V.  
 DT Journal  
 LA English  
 OS CASREACT 142:430102  
 GI




I

AB To develop novel PPAR $\gamma$  ligands, we synthesized thirteen 3-{4-(2-aminoethoxy)phenyl}propanoic acid derivs., which are designed based on the structures of rosiglitazone and 15d-PGJ2. Among these compds., compound I was found to be as potent as rosiglitazone in a binding assay and a preadipocyte differentiation test. Mol. modeling suggested that the nonyl group of I interacted with hydrophobic amino acid residues constructing the hydrophobic region of PPAR $\gamma$  protein where the alkyl chain of 15d-PGJ2 is expected to be located.

RE.CNT 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 3 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2004:394132 CAPLUS  
DN 141:391898  
TI Isolation and synthesis of isodihdropiperlonguminine  
AU Anuradha, V.; Srinivas, P. V.; Rao, J. Madhusudana  
CS Natural Products Laboratory, Organic Division-I, Indian Institute of Chemical Technology, Hyderabad, 500 007, India  
SO Natural Product Research (2004), 18(3), 247-251  
CODEN: NPRAAT; ISSN: 1478-6419  
PB Taylor & Francis Ltd.  
DT Journal  
LA English  
OS CASREACT 141:391898  
GI



I

AB The hexane extract of dried fruits of Piper longum on fractionation afforded a new alkamide, isodihdropiperlonguminine(I) and two Ph propanoic acid derivs. The structures of these compds. are established based on spectroscopic evidence and synthesis.

RE.CNT 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 4 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2004:226581 CAPLUS  
TI Novel dual PPAR  $\alpha$  and  $\gamma$  agonists derived from 2-alkoxy-3-phenyl propanoic acid series, which ameliorates metabolic abnormalities and reduces body weight  
AU Madhavan, G. R.; Chakrabarti, Ranjan; Reddy, Kalusam Anantha; Rajesh, B. M.; Rao, K. V. L. Narasimha; Rao, P. Bheema; Kumar, T. Ranjith;

Rajagopalan, R.

CS Metabolic Disorder Project Group, Dr. Reddy's Laboratories - Discovery Research, Hyderabad, 500050, India

SO Abstracts of Papers, 227th ACS National Meeting, Anaheim, CA, United States, March 28-April 1, 2004 (2004), MEDI-223 Publisher: American Chemical Society, Washington, D. C.

CODEN: 69FGKM

DT Conference; Meeting Abstract

LA English

AB Obesity is a disorder of fat accumulation and is associated with several risk factors known as metabolic syndrome. Present therapeutic approach to obesity is therefore also focused on overall management of metabolic syndrome. Peroxisome proliferator activated receptor (PPAR) is a member of nuclear receptor super family. Two of its isoforms - PPAR $\alpha$  and PPAR $\gamma$  are involved in the regulation of fat and carbohydrate metabolism and targets for hypolipidemic fibrates and antidiabetic thiazolidinediones. Considering the role of PPAR- $\alpha$  in catabolism of fat, we have initiated a program to discover a dual PPAR $\alpha$  /  $\gamma$ -agonist with greater specificity towards PPAR- $\alpha$ , so that it can be used for improving metabolic syndromes and body weight gain. We have investigated 1, 3-Benzoxazine-4(3H)-one derivs. of 2-alkoxy-3-Ph propanoic acid derivs. The SAR contains Cl, NO<sub>2</sub>, di-iso-Pr derivs. on aromatic ring of 1, 3-benzoxazinone and attachment of linker to -N' or -C' of the ring. Many compds. showed insulin sensitization and lipid lowering properties. DRF-2655 was selected as the lead mol. and resolved the racemic mixture in to its enantiomers (R and S). To our surprise both the enantiomers were having similar efficacy. Interestingly, DRF-2655 showed a significant body weight reduction in obese animal models along with good insulin sensitization and lipid lowering activity.

L1 ANSWER 5 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2004:226580 CAPLUS

TI New 2-ethoxy-3-(phenyl)propanoic acid derivatives as PPAR ligands

AU Bhuniya, Debnath; Iqbal, Javed; Chakrabarti, Ranjan; Mohan, Sankar; Narayanan, Sanju; Kumar, T. Ranjit; Suryaprakash, Raichur

CS Metabolic Disorder Group, Dr. Reddy-s Laboratories Ltd.- Discovery Research, Hyderabad, 500 049, India

SO Abstracts of Papers, 227th ACS National Meeting, Anaheim, CA, United States, March 28-April 1, 2004 (2004), MEDI-222 Publisher: American Chemical Society, Washington, D. C.

CODEN: 69FGKM

DT Conference; Meeting Abstract

LA English

AB Peroxisome proliferators-activated receptor (PPAR) is being recognized as versatile target for managing metabolic syndromes. Glitazones, as PPAR- $\gamma$  agonist, and fibtares as PPAR- $\alpha$  agonists are in the market for treatment of insulin resistant type 2 diabetes and dyslipidemia resp. In order to address diabetes and dyslipidemia with a single mol., a PPAR- $\alpha$ / $\gamma$  dual acting compound has been conceptualized. We believe that a proper combination of  $\alpha$ / $\gamma$  character may lead to a compound with such characteristics, without any significant PPAR- $\gamma$  related side effect. Along that line we have been working on design, synthesis and biol. evaluation of a series of 2-ethoxy-3-(phenyl)propanoic acids where Ph ring is linked through various spacers to a heterocycle. Quinazolinone as a representative heterocycle a general structure 1 with different examples have been screened on cell based PPAR assay. Selected compds. have been tested on relevant mice and rat models for type 2 diabetes and for dyslipidemia finally to come up with a lead structure having significant glucose and lipid lowering activity. Detailed synthesis and SAR will be presented in the form of poster.

L1 ANSWER 6 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2003:796655 CAPLUS

DN 139:292053  
 TI Etherification process for the preparation of 2-ethoxy-3-[4-[2-(4-methanesulfonyloxyphenyl)ethoxy]phenyl]propanoic acid derivatives  
 IN Larsson, Maria  
 PA Astrazeneca AB, Swed.; Astrazeneca UK Limited  
 SO PCT Int. Appl., 9 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003082812	A2	20031009	WO 2003-GB1395	20030328
	WO 2003082812	A3	20040108		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2478650	A1	20031009	CA 2003-2478650	20030328
	AU 2003226523	A1	20031013	AU 2003-226523	20030328
	BR 2003008297	A	20041228	BR 2003-8297	20030328
	EP 1492764	A2	20050105	EP 2003-745340	20030328
	EP 1492764	B1	20060628		
	R:				
	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	JP 2005521725	T	20050721	JP 2003-580280	20030328
	CN 1646487	A	20050727	CN 2003-807685	20030328
	NZ 534989	A	20060428	NZ 2003-534989	20030328
	AT 331704	T	20060715	AT 2003-745340	20030328
	ZA 2004006589	A	20050921	ZA 2004-6589	20040818
	NO 2004004045	A	20041018	NO 2004-4045	20040924
	US 2005215808	A1	20050929	US 2005-509654	20050505
	HK 1071353	A1	20061208	HK 2005-104064	20050513
PRAI	SE 2002-1005	A	20020402		
	WO 2003-GB1395	W	20030328		
OS	CASREACT 139:292053; MARPAT 139:292053				
GI					


\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB An efficient industrial-scale process for the preparation of 2-ethoxy-3-[4-[2-(4-methanesulfonyloxyphenyl)ethoxy]phenyl]propanoic acid derivs. [I; R = H, acid-protecting group; 1-(S)-2-ethoxy-3-[4-[2-(4-methanesulfonyloxyphenyl)ethoxy]phenyl]propanoic acid] is described which comprises the etherification of 2-ethoxy-3-(4-hydroxyphenyl)propanoate derivs. [II; e.g., Et (S)-2-ethoxy-3-(4-hydroxyphenyl)propanoate] with 2-(4-methanesulfonyloxyphenyl)ethyl derivs. [III; X = leaving group; e.g., 2-(4-methanesulfonyloxyphenyl)ethyl methanesulfonate] in the presence of a base (e.g., sodium carbonate) and using water as a diluent.

L1 ANSWER 7 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 1994:323168 CAPLUS  
 DN 120:323168  
 TI Dibasic (Amidinoaryl)propanoic Acid Derivatives as Novel Blood Coagulation

# Factor Xa Inhibitors

AU Nagahara, Takayasu; Yokoyama, Yukio; Inamura, Kazue; Katakura, Shin-ichi;  
 Komoriya, Satoshi; Yamaguchi, Hitoshi; Hara, Tsuyoshi; Iwamoto, Masahiro  
 CS Research Institute, Daiichi Pharmaceutical Company Ltd., Tokyo, 134, Japan  
 SO Journal of Medicinal Chemistry (1994), 37(8), 1200-7  
 CODEN: JMCMAR; ISSN: 0022-2623  
 DT Journal  
 LA English  
 GI



AB Since activated factor X (FXa) is a coagulant enzyme that generates thrombin and participates in both intrinsic and extrinsic coagulation pathways, inhibition of FXa may be more effective than inactivation of thrombin for interrupting blood coagulation. To assess the possible effectiveness of FXa inhibition as an anticoagulant, the authors designed and synthesized 3-(amidinoaryl)-2-[4-[(3S)-3-pyrrolidinyloxy]phenyl]propanoic acid derivs. as low mol. weight, nonpeptidic, orally active FXa inhibitors. These derivs. exhibited potent and highly selective anti-FXa activity in vitro and anticoagulant activity on oral administration. The most promising compound, I, inhibited 50% of FXa activity (IC50) at 0.07  $\mu$ M, doubled plasma recalcification time at 0.5  $\mu$ M, and significantly prolonged activated partial thromboplastin time at 100 mg/kg orally. In contrast with FXa inhibition, I showed no activity against thrombin (IC50 > 2000  $\mu$ M).

=> s human peroxisome proliferator-activated receptor alpha (PPAR)  
 MISSING OPERATOR 'ALPHA (PPAR'  
 The search profile that was entered contains terms or nested terms that are not separated by a logical operator.

=> s PPAR  
 8635 PPAR  
 1135 PPARS  
 L2 8768 PPAR  
 (PPAR OR PPARS)

=> s substituted phenylpropanoic acid derivatives  
 492689 SUBSTITUTED  
 1 SUBSTITUTEDS  
 492690 SUBSTITUTED  
 (SUBSTITUTED OR SUBSTITUTEDS)  
 821 PHENYLPROPANOIC  
 4311309 ACID  
 1568117 ACIDS  
 4812460 ACID  
 (ACID OR ACIDS)  
 340439 DERIVATIVES  
 1134482 DERIVS  
 1240054 DERIVATIVES  
 (DERIVATIVES OR DERIVS)  
 L3 8 SUBSTITUTED PHENYLPROPANOIC ACID DERIVATIVES  
 (SUBSTITUTED (W) PHENYLPROPANOIC (W) ACID (W) DERIVATIVES)

=> s L2 and L3

L4 7 L2 AND L3

=> d L4 1-7 bib abs

L4 ANSWER 1 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2006:1170541 CAPLUS

DN 146:19381

TI Design, synthesis, and evaluation of a novel series of  $\alpha$ -substituted phenylpropanoic acid derivatives as human peroxisome proliferator-activated receptor (PPAR)  $\alpha/\delta$  dual agonists for the treatment of metabolic syndrome

AU Kasuga, Jun-ichi; Yamasaki, Daisuke; Araya, Yoko; Nakagawa, Aya; Makishima, Makoto; Doi, Takefumi; Hashimoto, Yuichi; Miyachi, Hiroyuki

CS Institute of Molecular and Cellular Biosciences, University of Tokyo, Bunkyo-ku, Tokyo, 113-0032, Japan

SO Bioorganic & Medicinal Chemistry (2006), 14(24), 8405-8414  
CODEN: BMECEP; ISSN: 0968-0896

PB Elsevier Ltd.

DT Journal

LA English

AB A series of  $\alpha$ -alkyl-substituted phenylpropanoic acids was prepared as dual agonists of peroxisome proliferator-activated receptors alpha and delta (PPAR.alpha./ $\delta$ ). Structure-activity relationship studies indicated that the shape of the linking group and the shape of the substituent at the distal benzene ring play key roles in determining the potency

and the selectivity of PPAR subtype transactivation.

Structure-activity relationships among the amide series (10) and the reversed amide series (13) are similar, but not identical, especially in the case of the compds. bearing a bulky hydrophobic substituent at the distal benzene ring, indicating that the hydrophobic tail part of the mols. in these two series binds at somewhat different positions in the large binding pocket of PPAR.  $\alpha$ -Alkyl-substituted phenylpropanoic acids of (S)-configuration were identified as potent human PPAR.alpha./ $\delta$  dual agonists. Representative compds. exhibited marked nuclear receptor selectivity for PPAR.alpha. and PPAR.delta.. Subtype-selective PPAR activation was also examined by anal. of the mRNA expression of PPAR-regulated genes.

RE.CNT 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2005:1341960 CAPLUS

DN 144:232775

TI Design and synthesis of substituted phenylpropanoic acid derivatives as human peroxisome proliferator-activated receptor  $\alpha/\delta$  dual agonists

AU Kasuga, Jun-Ichi; Makishima, Makoto; Hashimoto, Yuichi; Miyachi, Hiroyuki

CS Institute of Molecular and Cellular Biosciences, University of Tokyo, Tokyo, 113-0032, Japan

SO Bioorganic & Medicinal Chemistry Letters (2006), 16(3), 554-558  
CODEN: BMCLE8; ISSN: 0960-894X

PB Elsevier B.V.

DT Journal

LA English

AB A series of phenylpropanoic acids was prepared as candidate dual agonists of peroxisome proliferator-activated receptors (PPAR)  $\alpha$  and  $\delta$ . Structure-activity relationship studies indicated that the shape of the linker moiety and the nature of the substituent at the distal benzene ring play key roles in determining the potency and selectivity of PPAR subtype transactivation. Optically active



RE.CNT 12        THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD  
                 ALL CITATIONS AVAILABLE IN THE RE FORMAT

CCCCCN(Cc1ccncc1)CCOCc1ccc(CCC(=O)O)cc1

I

RE.CNT 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4	ANSWER 4 OF 7	CAPLUS	COPYRIGHT 2007 ACS on STN
AN	2003:524282	CAPLUS	
DN	139:224182		
TI	Design, Synthesis, and Evaluation of Substituted Phenylpropanoic Acid Derivatives as Human Peroxisome Proliferator Activated Receptor Activators. Discovery of Potent and Human Peroxisome Proliferator Activated Receptor $\alpha$ Subtype-Selective Activators		
AU	Nomura, Masahiro; Tanase, Takahiro; Ide, Tomohiro; Tsunoda, Masaki; Suzuki, Masahiro; Uchiki, Hideharu; Murakami, Koji; Miyachi, Hiroyuki		
CS	Discovery Research Laboratories, Kyorin Pharmaceutical Co. Ltd., Nogi-chi, Shimotsugo-gun, Tochigi, 329-0114, Japan		
SO	Journal of Medicinal Chemistry (2003), 46(17), 3581-3599 CODEN: JMCMAR; ISSN: 0022-2623		
PB	American Chemical Society		
DT	Journal		
LA	English		
OS	CASREACT 139:224182		
GI			



AB Substituted phenylpropanoic acid  
 derivs. such as I are prepared as selective human peroxisome  
 proliferator activated receptor  $\alpha$  (PPAR. $\alpha$ .)  
 activators. Structure-activity relationships for the binding of a variety  
 of substituted phenylpropanoic acid  
 derivs. to human peroxisome proliferator activated receptors are  
 determined. The nature and the stereochem. of the substituent at the  
 $\alpha$ -position of the head part containing the carboxyl group, the distance  
 between the carboxyl group and the central benzene ring, the linking group  
 between the central benzene ring and the distal benzene ring, and the  
 substituent at the distal hydrophobic tail part of the mol. all play key  
 roles in determining the potency and selectivity of PPAR subtype  
 transactivation. Mol. mechanics calcns. of the conformers of  
 phenylpropanoic acid derivs. and of the enantiomers of an  
 $\alpha$ -ethyl-substituted phenylpropanoic acid derivative are discussed. I is  
 a particularly effective PPAR. $\alpha$ . activator with significant  
 selectivity for PPAR. $\alpha$ .. In rats, I decreases serum  
 cholesterol and lipids over five days of administration in a  
 dose-dependent manner and with a significantly greater efficacy than a  
 representative fibrate (bezafibrate) used for comparison. Phenylpropanoic  
 acid II is found to be a dual activator of PPAR. $\alpha$ . and of  
 PPAR. $\delta$ ..

RE.CNT 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 5 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 2003:184227 CAPLUS  
 TI Design, synthesis, and evaluation of substituted  
 phenylpropanoic acid derivatives as human  
 peroxisome proliferator-activated receptor activators: The discovery of  
 potent and human PPAR subtype-selective activators  
 AU Miyachi, Hiroyuki; Nomura, Masahiro; Tanase, Takahiro; Ide, Tomohiro;  
 Tsunoda, Masaki; Suzuki, Masahiro; Nagasawa, Michiaki; Uchiki, Hideharu;  
 Murakami, Koji  
 CS Discovery Research Laboratories, Kyorin Pharmaceutical Co., Ltd, Tochigi,  
 329-0114, Japan  
 SO Abstracts of Papers, 225th ACS National Meeting, New Orleans, LA, United  
 States, March 23-27, 2003 (2003), MEDI-261 Publisher: American Chemical  
 Society, Washington, D. C.  
 CODEN: 69DSA4  
 DT Conference; Meeting Abstract  
 LA English  
 AB A series of phenylpropanoic acids was prepared as part of a search for  
 subtype-selective PPAR  $\alpha$  activators. SAR studies indicated



that the nature of the substituent at the alpha position of the head part containing the carboxyl group, the distance between the carboxyl group and the central benzene ring, the linking group between the central benzene ring and the distal benzene ring, and the substituent at the distal hydrophobic tail of the mol. all play key roles in determining the potency and the selectivity. Transactivation study using chimeric PPAR alpha indicated that species-selective PPAR alpha transactivation was mediated via the interaction between the activator and the side chain of a crucial amino acid located in the helix three region of PPAR alpha. This study has led to the identification of potent and human PPAR alpha-selective derivs., which will be useful as candidate drugs for the treatment of metabolic disorders.

L4 ANSWER 6 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 2001:900080 CAPLUS  
 DN 136:318816  
 TI Design, synthesis and evaluation of substituted  
 phenylpropanoic acid derivatives as peroxisome  
 proliferator-activated receptor (PPAR) activators: novel human  
 PPAR.alpha.-selective activators  
 AU Miyachi, Hiroyuki; Nomura, Masahiro; Tanase, Takahiro; Takahashi, Yukie;  
 Ide, Tomohiro; Tsunoda, Masaki; Murakami, Koji; Awano, Katsuya  
 CS Kyorin Pharmaceutical Co., Ltd., Discovery Research Laboratories, Tochigi,  
 Shimotsuga-gun, Nogi-machi, 329-0114, Japan  
 SO Bioorganic & Medicinal Chemistry Letters (2001), Volume Date 2002, 12(1),  
 77-80  
 CODEN: BMCLE8; ISSN: 0960-894X  
 PB Elsevier Science Ltd.  
 DT Journal  
 LA English  
 OS CASREACT 136:318816  
 AB A series of substituted phenylpropanoic acid  
 derivs. was prepared as part of a search for subtype-selective human  
 peroxisome proliferator-activated receptor (PPAR) activators.  
 Structure-activity relationship studies indicated that the substituent at  
 the  $\alpha$ -position of the carboxyl group plays a key role in determining the  
 potency and the selectivity for PPAR transactivation.  
 RE.CNT 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 7 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 2001:284304 CAPLUS  
 DN 135:57906  
 TI Fluorine-Substituted Ligands for the Peroxisome Proliferator-Activated  
 Receptor Gamma (PPAR.gamma.): Potential Imaging Agents for  
 Metastatic Tumors  
 AU Kim, Sung-Hoon; Jonson, Stephanie D.; Welch, Michael J.; Katzenellenbogen,  
 John A.  
 CS Department of Chemistry, University of Illinois, Urbana, IL, 61801, USA  
 SO Bioconjugate Chemistry (2001), 12(3), 439-450  
 CODEN: BCCHES; ISSN: 1043-1802  
 PB American Chemical Society  
 DT Journal  
 LA English  
 AB The peroxisome proliferator-activated receptor gamma (PPAR  
 $\gamma$ ), a primary regulator of lipid metabolism, is present in many tumor  
 cell lines and animal tumor systems and, in some cases, can mediate  
 effective antitumor therapy with potent synthetic ligands. In an approach  
 to image tumors with positron-emission tomog. (PET) based on their content  
 of PPAR.gamma., we have synthesized two fluorine-substituted  
 analogs of a high affinity ligand from the phenylpropanoic acid class.  
 The analog having the highest affinity for PPAR.gamma. was  
 labeled with the positron-emitting radionuclide fluorine-18. In tissue  
 distribution studies in normal rats and in SCID mice bearing human breast  
 tumor xenografts, this compound did not show evidence of receptor-mediated

uptake. The prospects for using PPAR.gamma. as a target for imaging tumors may be limited by the low receptor concns. in tumors and by the pharmacokinetic behavior of this class of ligands, which appears to be more favorable for therapy than for imaging.

RE.CNT 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	60.53	60.74
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-10.92	-10.92

STN INTERNATIONAL LOGOFF AT 14:27:39 ON 09 FEB 2007

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssptalxn1621

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2		"Ask CAS" for self-help around the clock
NEWS	3	OCT 23	The Derwent World Patents Index suite of databases on STN has been enhanced and reloaded
NEWS	4	OCT 30	CHEMLIST enhanced with new search and display field
NEWS	5	NOV 03	JAPIO enhanced with IPC 8 features and functionality
NEWS	6	NOV 10	CA/Caplus F-Term thesaurus enhanced
NEWS	7	NOV 10	STN Express with Discover! free maintenance release Version 8.01c now available
NEWS	8	NOV 20	CA/Caplus to MARPAT accession number crossover limit increased to 50,000
NEWS	9	DEC 01	CAS REGISTRY updated with new ambiguity codes
NEWS	10	DEC 11	CAS REGISTRY chemical nomenclature enhanced
NEWS	11	DEC 14	WPIDS/WPINDEX/WPIX manual codes updated
NEWS	12	DEC 14	GBFULL and FRFULL enhanced with IPC 8 features and functionality
NEWS	13	DEC 18	CA/Caplus pre-1967 chemical substance index entries enhanced with preparation role

NEWS 14 DEC 18 CA/Capplus patent kind codes updated  
 NEWS 15 DEC 18 MARPAT to CA/Capplus accession number crossover limit increased to 50,000  
 NEWS 16 DEC 18 MEDLINE updated in preparation for 2007 reload  
 NEWS 17 DEC 27 CA/Capplus enhanced with more pre-1907 records  
 NEWS 18 JAN 08 CHEMLIST enhanced with New Zealand Inventory of Chemicals  
 NEWS 19 JAN 16 CA/Capplus Company Name Thesaurus enhanced and reloaded  
 NEWS 20 JAN 16 IPC version 2007.01 thesaurus available on STN  
 NEWS 21 JAN 16 WPIDS/WPINDEX/WPIX enhanced with IPC 8 reclassification data  
 NEWS 22 JAN 22 CA/Capplus updated with revised CAS roles  
 NEWS 23 JAN 22 CA/Capplus enhanced with patent applications from India  
 NEWS 24 JAN 29 PHAR reloaded with new search and display fields  
 NEWS 25 JAN 29 CAS Registry Number crossover limit increased to 300,000 in multiple databases

NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
 NEWS LOGIN Welcome Banner and News Items  
 NEWS IPC8 For general information regarding STN implementation of IPC 8  
 NEWS X25 X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 15:42:02 ON 09 FEB 2007

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 15:42:20 ON 09 FEB 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 8 FEB 2007 HIGHEST RN 920112-67-0  
 DICTIONARY FILE UPDATES: 8 FEB 2007 HIGHEST RN 920112-67-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 30, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=> ....Testing the current file.... screen


ENTER SCREEN EXPRESSION OR (END):end

=> screen 1006

L1 SCREEN CREATED

=>

Uploading C:\Program Files\Stnexp\Queries\10560746A.str



chain nodes :

1 2 3 4 5 12 13 14 15 16 17 18 19 20

ring nodes :

6 7 8 9 10 11

chain bonds :

1-2 1-20 2-3 2-5 3-4 4-6 9-12 12-13 13-14 13-17 14-15 14-16 17-18  
18-19

ring bonds :

6-7 6-11 7-8 8-9 9-10 10-11

exact/norm bonds :

1-2 1-20 2-5 4-6 13-17 14-15 14-16

exact bonds :

2-3 3-4 9-12 12-13 13-14 17-18 18-19

normalized bonds :

6-7 6-11 7-8 8-9 9-10 10-11

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:CLASS 7:CLASS 8:CLASS 9:CLASS

10:CLASS 11:CLASS 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom

19:Atom 20:Atom

L2 STRUCTURE UPLOADED


=> que L2 AND L1

L3 QUE L2 AND L1

=> d L2

L2 HAS NO ANSWERS

L2 STR



Structure attributes must be viewed using STN Express query preparation.

=> s L2 full

FULL SEARCH INITIATED 15:42:54 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 4810 TO ITERATE

100.0% PROCESSED 4810 ITERATIONS  
SEARCH TIME: 00.00.01

123 ANSWERS

L4 123 SEA SSS FUL L2

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE  
ENTRY

TOTAL  
SESSION

FULL ESTIMATED COST

172.10

172.31

FILE 'CAPLUS' ENTERED AT 15:43:01 ON 09 FEB 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 9 Feb 2007 VOL 146 ISS 8

FILE LAST UPDATED: 8 Feb 2007 (20070208/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s L2

REGISTRY INITIATED  
Substance data SEARCH and crossover from CAS REGISTRY in progress...  
Use DISPLAY HITSTR (or FHITSTR) to directly view retrieved structures.

SAMPLE SEARCH INITIATED 15:43:06 FILE 'REGISTRY'  
SAMPLE SCREEN SEARCH COMPLETED - 255 TO ITERATE

100.0% PROCESSED 255 ITERATIONS 5 ANSWERS  
SEARCH TIME: 00.00.02

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*  
PROJECTED ITERATIONS: 4142 TO 6058  
PROJECTED ANSWERS: 5 TO 234



L5 5 SEA SSS SAM L2

L6 5 L5

=> d L6 1-5 bib abs

L6 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2007:61504 CAPLUS  
DN 146:142376  
TI Preparation of phenylpropionic acid derivatives and pharmaceutical  
compositions thereof  
IN Bjoerk, Seth  
PA Astrazeneca AB, Swed.  
SO PCT Int: Appl., 57pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2007008156	A1	20070118	WO 2006-SE864	20060710
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,				
	CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,				
	GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,				
	KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN,				
	MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU,				
	SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG,				
	US, UZ, VC, VN, ZA, ZM, ZW				
	RW:				
	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,				
	IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ,				
	CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH,				
	GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,				
	KG, KZ, MD, RU, TJ, TM				
PRAI	SE 2005-1644	A	20050711		
GI					



AB The title phenylpropionic acid derivs. I [wherein n = 1-2; R1 = H, Cl, CF3, or OCF3; R2 = H or F; R3 = alkyl] or tert-butylamine salts thereof were prepared as PPAR active compds. for treatment of metabolic syndrome including type 2 diabetes mellitus (no data). For example, II and II•tert-butylamine were prepared in a multi-step synthesis. Pharmaceutical compns. were described.


RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2005:1335635 CAPLUS  
DN 144:69628  
TI Preparation of phenoxyacetamide derivatives as modulators of peroxisome proliferator-activated receptors (PPAR)  
IN Alstermark, Eva-Lotte Lindstedt; Olsson, Anna Christina; Li, Lanna  
PA Swed.  
SO U.S. Pat. Appl. Publ., 47 pp., Cont.-in-part of U.S. Ser. No. 499,261.  
CODEN: USXXCO  
DT Patent  
LA English  
FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2005282822	A1	20051222	US 2004-26806	20041230
	WO 2003051821	A1	20030626	WO 2002-GB5738	20021218
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	WO 2003051822	A1	20030626	WO 2002-GB5744	20021218
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CN 1896045	A	20070117	CN 2006-10007173	20021218

WO 2004056748	A1	20040708	WO 2003-GB5602	20031219
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
WO 2004113270	A2	20041229	WO 2004-EP6597	20040617
WO 2004113270	A3	20050331		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1676833	A1	20060705	EP 2006-5766	20040617
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR			
JP 2005336209	A	20051208	JP 2005-235794	20050816
JP 2006045240	A	20060216	JP 2005-253346	20050901
JP 2006298924	A	20061102	JP 2006-123399	20060427
JP 2006298925	A	20061102	JP 2006-139673	20060519
PRAI SE 2001-4334	A	20011219		
WO 2002-GB5738	W	20021218		
WO 2002-GB5744	A	20021218		
GB 2002-29931	A	20021221		
GB 2003-14079	A	20030618		
WO 2003-GB305602	A	20031219		
WO 2004-EP6597	A	20040617		
US 2005-499261	A2	20050304		
CN 2002-828123	A3	20021218		
JP 2003-552709	A3	20021218		
JP 2003-552710	A3	20021218		
JP 2004-561668	A3	20031219		
EP 2004-740044	A3	20040617		
JP 2006-515989	A3	20040617		

OS  
GI



AB The phenyl-, phenoxy-, or phenylthioalkanamidetitle compds., (in particular phenoxyacetamide derivs.) (I) [A is situated in the ortho, meta or para position and represents CR3R4CR1R2COR, CR3:CR1COR (wherein R = H, alkyl, (un)substituted HO or NH2; R1 = alkyl, aryl, alkenyl, alkynyl, or when A is CR3R4CR1R2COR, R1 can also be cyano, (un)substituted HO, SH, OCONH2, SO2NH2, CO2H, etc.; R2 = H, halogen, alkyl, aryl, alkylaryl; R3, R4 = H, alkyl, aryl, alkylaryl); Y = O, S, a single bond; n = an integer



of 1-4; X = alkyl; R5, R6 = H, each (un)substituted C1-13 alkyl, C2-10 alkenyl, or C2-10 alkynyl; or R5, R6 = each (un)substituted C3-8 cycloalkyl, C3-C8 cycloalkenyl, aryl, heterocyclyl, or heteroaryl; or R5 and R6 together with the nitrogen atom to which they are attached form a single or a fused heterocyclic system] are prepared These compds. are useful in treating clin. conditions including lipid disorders (dyslipidemias) whether or not associated with insulin resistance, and other manifestations of the metabolic syndrome. Thus, a solution of 0.598 g N-butyl-N-[2-fluoro-4-(trifluoromethyl)benzyl]amine and 0.593 g [4-((2S)-2,3-diethoxy-3-oxopropyl)phenoxy]acetic acid in 20 mL CH<sub>2</sub>Cl<sub>2</sub> was treated with 0.80 mL N,N-diisopropylethylamine and 0.674 g O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate and the reaction mixture was stirred at room temperature overnight to give, after workup and silica gel chromatog., 74% Et (2S)-3-[4-[2-[butyl[2-fluoro-4-(trifluoromethyl)benzyl]amino]-2-oxoethoxy]phenyl]-2-ethoxypropanoate (II). A solution of 0.748 g II in 70 mL MeCN was treated with 35 mL 0.10 M LiOH and the reaction mixture was stirred at room temperature overnight, neutralized with 5% HCl, concentrated, acidified with 5% HCl, and extracted



with

EtOAc to give 97% (2S)-3-[4-[2-[butyl[2-fluoro-4-(trifluoromethyl)benzyl]amino]-2-oxoethoxy]phenyl]-2-ethoxypropanoic acid (III). III showed EC<sub>50</sub> of 0.001 µmol/L for human PPAR $\alpha$ .

L6 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2007 ACS on STN  
 AN 2004:1154649 CAPLUS  
 DN 142:93514  
 TI Preparation of phenylpropanoic acid derivatives as PPAR $\alpha$  agonists  
 IN Li, Lanna; Lindstedt-Alstermark, Eva-Lotte; Olsson, Christina  
 PA Astrazeneca Ab, Swed.  
 SO PCT Int. Appl., 100 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004113270	A2	20041229	WO 2004-EP6597	20040617
	WO 2004113270	A3	20050331		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, VZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2004249409	A1	20041229	AU 2004-249409	20040617
	CA 2528234	A1	20041229	CA 2004-2528234	20040617
	US 2005148656	A1	20050707	US 2003-518777	20040617
	EP 1675820	A2	20060705	EP 2004-740044	20040617
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
	EP 1676833	A1	20060705	EP 2006-5766	20040617
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
	BR 2004011484	A	20060725	BR 2004-11484	20040617
	CN 1835913	A	20060920	CN 2004-80023304	20040617
	JP 2006527730	T	20061207	JP 2006-515989	20040617
	US 2005282822	A1	20051222	US 2004-26806	20041230
	NO 2005005892	A	20060106	NO 2005-5892	20051212
	JP 2006298925	A	20061102	JP 2006-139673	20060519
	US 2006258866	A1	20061116	US 2006-477168	20060628

PRAI	GB 2003-14079	A	20030618
	SE 2001-4334	A	20011219
	WO 2002-GB5738	W	20021218
	WO 2002-GB5744	A	20021218
	GB 2002-29931	A	20021221
	WO 2003-GB305602	A	20031219
	EP 2004-740044	A3	20040617
	JP 2006-515989	A3	20040617
	WO 2004-EP6597	W	20040617
	US 2005-518777	A3	20050303
	US 2005-499261	A2	20050304
OS	MARPAT 142:93514		
GI			





AB Title compds. represented by the formula I [wherein A = CR<sub>3</sub>(R<sub>4</sub>)CR<sub>1</sub>(R<sub>2</sub>)COR or C(R<sub>3</sub>):C(R<sub>1</sub>)COR; R = H, alkoxy, (alkyl)aryloxy, amino, etc.; R<sub>1</sub> = alkyl, aryl, alkenyl, alkynyl, etc.; R<sub>2</sub> = H, halo, alkyl, (alkyl)aryl; R<sub>3</sub>, R<sub>4</sub> = independently H, alkyl, (alkyl)aryl; T = O, S or a single bond; n = 1-4; R<sub>5</sub>, R<sub>6</sub> = independently selected substituent comprising C, H, N, O, S, Se, P or halo; with provisos; optical isomers and racemates thereof as well as pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof] were prepared as PPAR $\alpha$  agonists. For example, II was given in a multi-step synthesis starting from the reaction of 2,4-difluorobenzylamine with octanoic acid. I had EC<sub>50</sub> values of less than 0.1  $\mu$ mol/L for PPAR $\alpha$  and showed the ration of the EC<sub>50</sub>(PPAR $\gamma$ ) with EC<sub>50</sub>(PPAR $\alpha$ ) is greater than 150:1. Thus, I and their pharmaceutical compns. are useful for the treatment of clin. conditions including lipid disorders (dyslipidemias) whether or not associated with insulin resistance (no data).

L6 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2004:2837 CAPLUS  
DN 140:59411  
TI Preparation of phenoxyalkanamides as amide linker peroxisome proliferator activated receptor agonists for treating and/or preventing diabetes mellitus and syndrome X  
IN Ferritto Crespo, Rafael; Martin, Jose Alfredo; Martin-Ortega, Finger Maria Dolores; Rojo Garcia, Isabel; Shen, Quanrong; Warshawsky, Alan M.; Xu, Yanping  
PA Eli Lilly and Company, USA  
SO PCT Int. Appl., 168 pp.  
CODEN: PIXXD2  
DT Patent  
LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE	
PI	WO 2004000789	A1	20031231	WO 2003-US16207	20030611	
	WO 2004000789	A9	20040311			
	W:			AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW		
	RW:			GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG		
	CA 2488972	A1	20031231	CA 2003-2488972	20030611	
	AU 2003241579	A1	20040106	AU 2003-241579	20030611	
	EP 1517882	A1	20050330	EP 2003-731326	20030611	
	R:			AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK		
	BR 2003011834	A	20050412	BR 2003-11834	20030611	
	CN 1662487	A	20050831	CN 2003-814173	20030611	
	JP 2005529975	T	20051006	JP 2004-515700	20030611	
	US 2006111406	A1	20060525	US 2004-517581	20041208	
PRAI	US 2002-390102P	P	20020619			
	WO 2003-US16207	W	20030611			
OS	MARPAT 140:59411					
GI						



AB The present invention is directed to phenoxyalkanamides (shown as I; variables defined below; e.g. II), compns., and their use as peroxisome proliferator activated receptor agonists for treating and/or preventing diabetes mellitus and syndrome X. The binding and cotransfection efficacy values found for compds. of this invention that are useful for modulating a PPAR $\alpha$  receptor are about <100 nM and >50%, resp. Although the methods of preparation are not claimed, apprx.140 example prepsns. of I are included. For example, II was prepared in 3 steps starting from (2S)-2-ethoxy-3-(4-hydroxyphenyl)propionic acid Me ester, (2S)-2-hydroxypropionic acid benzyl ester and involving intermediates (2S)-3-[4-[[[(1R)-1-[(benzyloxy)carbonyl]ethyl]oxy]phenyl]-2-ethoxypropionic acid Et ester and (2S)-3-[4-[[[(1R)-1-carboxyethyl]oxy]phenyl]-2-ethoxypropionic acid. For I: R1 = H, C1-C8 alkyl, C3-C6 cycloalkyl, aryl-C0-4-alkyl, heteroaryl-C0-4-alkyl, aminoC1-C4alkyl, C3-C6 cycloalkylaryl-C0-2-alkyl, arylheteroC1-C8alkyl, -CHC(O)C1-C4 alkoxy, C0-4-alkyl-C(O)heteroC1-C8alkyl, and -CH2C(O)-R15R16. R2 = C1-C8 alkyl, C3-C6 cycloalkyl, aryl-C0-C4-alkyl, heteroaryl-C0-C4-alkyl, heteroC1-C6cycloalkylaryl, heteroC1-C6cycloalkylarylC1-C4alkyl,

aminoC1-C4alkyl, C3-C6 cycloalkylaryl-C0-C2-alkyl, arylheteroC1-C8alkyl, C0-C4-alkyl-C(O)heteroC1-C8alkyl, -CH(C(O)OCH3)benzyl, and -CH2C(O)R15''R16''. R1 and R2 together may form a heterocyclic ring which heterocyclic ring is (un)substituted with 1-3 substituents R1' and which heterocyclic ring is optionally fused with an aryl; E = C(R3)(R4)A, (CH2)nCOOR13, aryl-C0-C4-alkyl, thio-C1-C4-alkyl, thioaryl, arylC1-C4alkoxy, C1-C4alkoxy C1-C4alkyl, aminoaryl, and aminoC1-C4alkyl. R5 and R6 = H, C1-C8 alkyl, aryl-C0-C4-alkyl, heteroaryl-C0-C4-alkyl, C3-C6 cycloalkyl, aryl-C0-C2-alkyl, C3-C6 cycloalkyl-C0-2-alkyl, and -CH2C(O)R17R18.

RE.CNT 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2007 ACS on STN  
AN 2003:491168 CAPLUS  
DN 139:69049  
TI Preparation of substituted phenylpropionic acid derivatives as agonists to human peroxisome proliferator-activated receptor alpha (PPAR)  
IN Alstermark Lindstedt, Eva-Lotte; Olsson, Anna Christina; Li, Lanna  
PA Astrazeneca AB, Swed.; Astrazeneca UK Limited  
SO PCT Int. Appl., 40 pp.  
CODEN: PIXXD2


DT Patent  
LA English

FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003051821	A1	20030626	WO 2002-GB5738	20021218
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2470491	A1	20030626	CA 2002-2470491	20021218
	AU 2002366315	A1	20030630	AU 2002-366315	20021218
	EP 1458673	A1	20040922	EP 2002-804964	20021218
	EP 1458673	B1	20060906		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
	BR 2002014988	A	20041214	BR 2002-14988	20021218
	HU 200402133	A2	20050228	HU 2004-2133	20021218
	CN 1620422	A	20050525	CN 2002-828123	20021218
	CN 1620423	A	20050525	CN 2002-828155	20021218
	US 2005171204	A1	20050804	US 2003-499261	20021218
	JP 2005526011	T	20050902	JP 2003-552709	20021218
	JP 3784804	B2	20060614		
	TW 253444	B	20060421	TW 2002-91136518	20021218
	NZ 533276	A	20060428	NZ 2002-533276	20021218
	TW 255807	B	20060601	TW 2002-91136519	20021218
	AT 338743	T	20060915	AT 2002-804964	20021218
	CN 1896045	A	20070117	CN 2006-10007173	20021218
	ZA 2004004657	A	20050829	ZA 2004-4657	20040611
	ZA 2004004658	A	20060222	ZA 2004-4658	20040611
	NO 2004003023	A	20040715	NO 2004-3023	20040715
	US 2005282822	A1	20051222	US 2004-26806	20041230
	JP 2005336209	A	20051208	JP 2005-235794	20050816
	JP 2006298924	A	20061102	JP 2006-123399	20060427
PRAI	SE 2001-4334	A	20011219		
	CN 2002-828123	A3	20021218		
	JP 2003-552709	A3	20021218		

JP 2003-552710	A3	20021218
WO 2002-GB5738	W	20021218
WO 2002-GB5744	A	20021218
GB 2002-29931	A	20021221
GB 2003-14079	A	20030618
WO 2003-GB305602	A	20031219
WO 2004-EP6597	A	20040617
US 2005-499261	A2	20050304

OS MARPAT 139:69049/  
GI



AB The S enantiomer of I, n = 1 or 2, (C<sub>6</sub>H<sub>13</sub> = hexyl) as well as their pharmaceutically acceptable salts, solvates, crystalline forms and prodrugs are synthesized using various solvents and in presence of charcoal-supported palladium catalyst. The utility of these compds. in clin. conditions such as lipid disorders (dyslipidemias) whether or not associated with insulin resistance and therapeutic and other pharmaceutical activities is also investigated. For example, (2S)-3-(4{2-[benzyl(hexyl)amino]-2-oxoethoxy}phenyl)2-ethoxypropionic acid was prepared in 58% yield via reaction of (2S)-2-ethoxy-3-(4-hydroxyphenyl)propanoate and benzyl bromoacetate.

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	15.09	188.32
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-3.90	-3.90

STN INTERNATIONAL LOGOFF AT 15:44:06 ON 09 FEB 2007